

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Invasive pathogens and pests may threaten a multipurpose tree species: the European chestnut as a case study

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/155486> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

This is an author version of the contribution published on:

Questa è la versione dell'autore dell'opera:

[*International Forestry Review*, 16(5), 2014]

ovvero [Gonthier P., Ferracini C., pagg. 335]

The definitive version is available at:

La versione definitiva è disponibile alla URL:

[*inserire* URL sito editoriale]

How invasive pathogens and pests may threaten a multipurpose tree species: the European chestnut as a case study

Paolo Gonthier and Chiara Ferracini

University of Torino, Dept. of Agricultural, Forest and Food Sciences, Via L. da Vinci 44, 10095 Grugliasco, Italy

Keywords: *Cryphonectria parasitica*, *Dryocosmus kuriphilus*, *Gnomoniopsis castanea*, impact, *Phytophthora*

The European chestnut (*Castanea sativa*) is a multipurpose tree species valued for nuts, timber, tannins, hydrogeological protection and landscape. It has long been serving as a substitute of cereals and as a main source of carbohydrates and firewood (fuel) for people living in mountain areas of southern Europe. In this paper we show how invasive pathogens and pests of chestnut accidentally introduced or of unknown origin have significantly impacted the rural economy, and we propose that tree decline currently observed in several coppices and orchards may be the result of interaction amongst them, possibly within the frame of changing socio-economic and climatic conditions. Ink disease caused by the oomycetes *Phytophthora* spp. and chestnut blight due to the fungal pathogen *Cryphonectria parasitica* spread epidemically since the mid-1800s and first half of 1900s, respectively, threatening the survival of the European chestnut. The recent introduction and invasion of the oriental chestnut gall wasp *Dryocosmus kuriphilus* and the newly described nut rot caused by the fungus *Gnomoniopsis castanea*, which may locally decrease nut production by 80%, are regarded as two important additional factors leading to the abandonment of chestnut stands. Actual social consequences under an economic, cultural and political perspective will be discussed.